

Indian Journal of Mathematics

Volume 37, No. 2, 1995

CONTENTS

Mathew He

ON THE ZEROS OF WEIGHTED FABER POLYNOMIALS 79-93

Abstract: Weighted Faber polynomials $\{F_n(z; g)\}$ associated with a domain E and a weight function $g(z)$ play a very important role in the study of the asymptotic properties of orthogonal polynomials in the complex domain. Here, we present a new determinant representation of $F_n(z; g)$ which relates the zeros of $F_n(z; g)$ to the eigenvalues of a certain matrix and study the location and the asymptotic distribution of the zeros of $F_n(z; g)$ mainly in dependence on the smoothness of the weight function and the boundary of the domain.

S. Fridli

MEAN CONVERGENCE OF WALSH-FOURIER SERIES 95-101

Abstract: It is known that the integrability of a function does not guarantee the convergence of the corresponding Walsh-Fourier series. An additional condition that implies the convergence can be made by the L^1 modulus of continuity-Dini-Lipschitz condition-, or by requiring that the function belongs to the narrower space, say $L^p[0, 1]$ ($1 < p \leq \infty$). Another possibility is to give a convergence condition with respect to the Walsh-Fourier coefficients. In this paper we formalize such a condition by means of a shifted Sidon type inequality for the Walsh-Dirichlet kernels and by using the concepts of dyadic Hardy space and generalized de la Vallée Poussin means.

Tai-Jan Huang and Young-Ye Huang

FIXED POINT THEOREMS FOR LEFT REVERSIBLE SEMIGROUPS IN
COMPACT MATRIC SPACES 103-105

Abstract: It is shown that a left reversible semigroups of contractive selfmaps on a compact metric self space (M, d) has a unique fixed point ξ , and for any x in M and any t in S the iterates $t^n x$ converges ξ .

Z. Govindarajulu

A NOTE ON TWO-STAGE FIXED-WIDTH INTERVAL ESTIMATION
PROCEDURE FOR NORMAL VARIANCE 107-112

Abstract: A large-sample two-stage solution is obtained for the problem of setting fixed-width confidence intervals for the normal variance. This method yields a substantial reduction in the second sample size required by other existing methods.

E. Karapinar and G. SH Guseinov

THE BOUNDEDNESS OF SOLUTIONS OF SECOND-ORDER
DIFFERENCE EQUATIONS

113-122

Abstract: In this note, some simple conditions for the boundedness of all solutions of a second-order difference equations on the half-line are given.

B. Mond and J. Pečarić

A SIMPLE PROOF OF GENERALIZED INEQUALITIES OF BHAGWAT
AND SUBRAMANIAN AND SOME CONVERSE RESULTS

123-128

Abstract: Bhagwat and Subramanian proved a number of inequalities involving powers of positive operators. Here a simple proof of generalization of these inequalities is given. Converses for some special cases are also established.

Sang Chul Lee and Byung Soo Lee

SOME GENERALIZATIONS OF MINIMAX INEQUALITY

129-135

Abstract: We obtain a generalized minimax inequality using H - KKM theorem.

Shih-Sen Chang and Yi-Hai Ma

KKM TECHNIQUE AND ITS APPLICATIONS

137-150

Abstract: In this paper, the Knaster-Kuratowski-Mazurkeiwicz technique (KKM technique, in short) is presented. Using this technique a new alternative theorem and a new coincidence theorem are established. The results obtained in this paper unify and generalize the corresponding results in the recent works [2,10,11,15,16].

Adrian Constantin

A RANDOM INTEGRAL EQUATION WITH APPLICATIONS

151-163

Abstract: We will investigate the existence, uniqueness and asymptotic behavior of the random solution for the stochastic integral equation

$$x(t; w) = h(t; w) + \int_0^t k(t, s; w) f(s, x(s; w)) ds, \quad t \geq 0.$$

and we give some examples of equations of this form which arise in hereditary mechanics and population growth modeling.

A. K. Nandakumaran and Raju K. George

PARTIAL EXACT CONTROLLABILITY OF A LINEAR THERMOELASTIC
SYSTEM

165-174

Abstract: In this article, we prove the partial exact controllability of a one dimensional linear thermoelastic system. We use $RHUM$ method which is a variation of HUM method to study the present system.
