

Indian Journal of Mathematics

Volume 44, No. 1, 2002

CONTENTS

Ingrid Carbone

TROTTER'S TYPE THEOREMS FOR GENERALIZED SZÁSZ-MIRAKJAN
OPERATORS IN POLYNOMIAL WEIGHTED FUNCTION SPACES

1-20

Abstract: We consider the class of degenerate elliptic second order differential operators $Au(x) := \alpha(x) \ddot{u}(x) (x \geq 0)$ defined in some subspaces of the polynomial weighted function spaces $W_N^0 (N \geq 2)$ of all continuous functions u on $[0, +\infty[$ such that $\frac{u(x)}{1+x^N}$ vanishes at infinity. We show that, under suitable hypothesis on the function α , these operators are the generators of C_0 -semigroups on W_N^0 , that can be represented as a limit of powers of discrete-type positive linear operators that have been introduced earlier and are constructed by means of the coefficient α . We also derive some regularity results about these semigroups and, as an application, the solutions of the abstract Cauchy problem related to them.

All these results extend the corresponding ones obtained recently in the space W_2^0 .

Shaun Cooper and Heung Yeung Lam

SUMS OF TWO, FOUR, SIX AND EIGHT SQUARES AND TRIANGULAR
NUMBERS: AN ELEMENTARY APPROACH

21-40

Abstract: An elementary and self contained method is given for determining formulas for the number of representations of an integer as a sum of two, four, six and eight squares or triangular numbers. Our method uses Ramanujan's $|\Psi|$ summation formula and is based on K . Venkatachaliengar's elementary approach to elliptic functions.

Nawneet Hooda and Babu Ram

CONVERGENCE OF CERTAIN MODIFIED COSINE SUM

41-46

Abstract: In this paper, we generalize the results of Kumari and Ram [4], and Teljakovskii [9] for a more general class of sequences.

M. Imdad and Q. H. Khan

A COMMON FIXED POINT THEOREM FOR SIX MAPPINGS SATISFYING A
RATIONAL INEQUALITY

47-57

Abstract: Using notions of compatibility, weak compatibility and commutativity, we prove a common fixed point theorem for six mappings satisfying a rational inequality which unifies and improves earlier fixed point theorems due to Fisher Kannan, Hardy-Rogers and others. A related example is also furnished.

Mohammed Imdad and Tariq Iqtadar KhanCOINCIDENCE AND COMMON FIXED POINTS OF NONLINEAR
HYBRID CONTRACTIONS

59-71

Abstract: Using certain weak conditions of commutativity some common fixed point theorems for single-valued mappings and multi-valued mappings satisfying a rational inequality for six mappings are obtained which generalize some earlier results due to Cho et al., Fisher, Diviccaro et al., V. Popa, and others.

M. K. R. S. Veera Kumar $g^\#$ -SEMI-CLOSED SETS IN TOPOLOGICAL SPACES

73-87

Abstract: In this paper, we introduce and study a new class of sets, namely $g^\#$ -semi-closed sets for topological spaces. Applying $g^\#$ -semi-closed sets, we introduce and study four new classes of spaces, namely ${}^\#T_b$ spaces, $T_b^\#$ spaces, $T_b^{\#\#}$ spaces and ${}_\alpha T_b^\#$ spaces. The class of $T_b^\#$ spaces (resp. ${}^\#T_b$ spaces) is the dual of the class of ${}^\#T_b$ spaces (resp. ${}_\alpha T_b^\#$ spaces) to the class of T_b spaces (resp. $T_{1/2}$ spaces). We also introduce and study $g^\#$ s -continuous and $g^\#$ s -irresolute maps.

P. N. Natarajan

CHARACTERIZATION OF SOME MATRIX CLASSES

89-94

Abstract: In this paper the matrix classes (bv, Q) and (bv_0, Q) are characterized.

A. A. M. Wasike

PERIODIC SOLUTIONS OF A SYSTEM OF DELAY DIFFERENTIAL EQUATIONS 95-117

Abstract: We use a planar analysis approach to show the existence of periodic solutions of the system of delay differential equations $x'(t) = dL[x(t-\tau) - x(t)] + f(x)$, for $d, \tau > 0$, L a 2×2 positive semi definite matrix, and f is a smooth nonlinear function satisfying some negative feedback conditions. This involves finding a cone \mathcal{K} in the phase space $C([- \tau, 0], R^2)$ such that every solution $x_t(d, \varphi)$ of the equations subject to $x(t)|_{[- \tau, 0]} = \varphi(t)$ with $\varphi \in \mathcal{K}$ returns to \mathcal{K} after some time $0 < T(d, \tau) < \infty$ under a certain completely continuous operator $\mathcal{A} : \varphi \mathcal{K} \rightarrow xT(d, \varphi)(d, \varphi) \in \mathcal{K}$. Then, we show that there is a $\varphi \in \mathcal{K}$ such that $\mathcal{A}\varphi = \varphi$. This fixed point will correspond to the periodic solution of the equation with $x(t)|_{[- \tau, 0]} = \varphi(t)$. However, $0 \in \mathcal{K}$ is also a fixed point corresponding to the trivial periodic solution. Thus, to show the existence of a nontrivial periodic solution, we show that zero is an elective point with respect to \mathcal{A} .
